


# Cointegration

(Hamilton: Chapters 19, Zivot and Wang: Chapter 12 )  
Isabel Casas

- Spurious regression
- Cointegration
  - Phillip's triangular representation
  - Error-correction representation
- Test for cointegration
  - When the cointegrating vector is pre-defined
  - Estimating the cointegrating vector


## Example: spurious regression

```
> epsilon1=rnorm(1000)
> epsilon2=rnorm(1000)
> y1=cumsum(epsilon1)
> y2=cumsum(epsilon2)
```



# Example: spurious regression

Are they linearly related?



# Example: spurious regression

Run the regression:  $y_{1t} = \beta y_{2t} + e_t$

```
> summary(model)
```

Call:

```
lm(formula = y1 ~ -1 + y2)
```

Residuals:

	Min	1Q	Median	3Q	Max
	-30.539	-14.433	-9.310	-2.760	6.783

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
y2	0.36370	0.01898	19.17	<2e-16 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 12.41 on 999 degrees of freedom


Multiple R-squared: 0.2689, Adjusted R-squared: 0.2681

F-statistic: 367.4 on 1 and 999 DF, p-value: < 2.2e-16

## Example: spurious regression

$$y_{1t} = 0.36y_{2t} + e_t \quad \hat{\sigma} = 12.41$$

- The slope coefficient is 0.36 and appears as highly significant
- The  $R^2 = 0.27$  is quite good and it gets better as the sample size increases
- The residuals autocorrelation:



## Example: spurious regression

### The Durbin-Watson test

$H_0$  : autocorrelation of the innovations is zero

```
> library(lmtest)
> dwtest(model)
```

Durbin-Watson test

```
data: model
```

```
DW = 0.0068, p-value < 2.2e-16
```

```
alternative hypothesis: true autocorrelation is greater than 0
```

- These are the properties of a spurious regression
- It happens because the two variables are  $I(1)$  and not *cointegrated*
- If we regress  $\Delta y_{1t}$  on  $\Delta y_{2t}$ , the correct relationship is revealed

## Example: spurious regression

```
> deltay1=diff(y1)
> deltay2=diff(y2)
> model2<-lm(deltay1~-1 + deltay2)
```

Call:

```
lm(formula = deltay1 ~ -1 + deltay2)
```

Residuals:

Min	1Q	Median	3Q	Max
-3.07848	-0.72668	0.00278	0.66122	3.06710

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
deltay2	0.006419	0.030439	0.211	0.833

Residual standard error: 0.957 on 998 degrees of freedom


Multiple R-squared: 4.456e-05, Adjusted R-squared: -0.0009574

F-statistic: 0.04447 on 1 and 998 DF, p-value: 0.833



## Example: spurious regression

- The residuals of the regression of the first difference are now serially uncorrelated
- However, we have lost information taking the first difference
- And, it might be difficult to interpret the results back on the real series  $y_{1t}, y_{2t}$



# Statistical implication of spurious regression

$$y_{1t} = \beta y_{2t} + e_t$$

- $y_{1t}$  and  $y_{2t}$  are  $I(1)$  variables
- Since they are not cointegrated, the true value of  $\beta = 0$
- $\hat{\beta}$  do not converge in probability to zero but instead converges in distribution to a non-normal random variable not necessarily centered at zero.
- The usual OLS t-statistics for testing  $\beta = 0$  diverge to  $\pm\infty$  as  $T \rightarrow \infty$
- The usual  $R^2$  converges to 1 as  $T \rightarrow \infty$
- Regression with  $I(1)$  variables only makes sense if they are cointegrated.

# Applications where this is important

Examples of financial series that depend of each other.

## Definition: cointegration

A vector time series  $\mathbf{y}_t = \{y_{1t}, \dots, y_{nt}\}'$  is said to be *cointegrated of order 1* if

- ❶ Each of the series taken individually are  $I(1)$  (they have a unit root) and,
  - ❷ There is a combination of the series  $\mathbf{a}'\mathbf{y}_t$  which is  $I(0)$  for some non zero  $\mathbf{a}$
- Although changes in the individual elements of  $\mathbf{y}_t$  drift apart, there is a long-run equilibrium relation trying to keep these components together (they cannot escape from each other)
  - $\mathbf{a}$  is called the *cointegrating vector*
  - Simile: A drunk and her dog (Murray, 1994)

Q: cointegrated of order  $n$ ?

## Definition: cointegration

- This linear relationship  $\alpha$  is also called the *atractor* or as Murray says an *error-correction mechanism*
- The two variables are allowed to diverge in the short-run,
- In the long-run they have to converge to a common region

## Definition: cointegration

- $y_{1t} \sim I(0) \Rightarrow a + by_{1t} \sim I(0)$
- $y_{1t} \sim I(1) \Rightarrow a + by_{1t} \sim I(1)$
- $y_{1t}, y_{2t} \sim I(0) \Rightarrow ay_{1t} + by_{2t} \sim I(0)$
- $y_{1t} \sim I(0), y_{2t} \sim I(1) \Rightarrow ay_{1t} + by_{2t} \sim I(1)$
- 

$$y_{1t}, y_{2t} \sim I(1) \Rightarrow \begin{cases} ay_{1t} + by_{2t} \sim I(1) & \text{in general} \\ ay_{1t} + by_{2t} \sim I(0) & \text{if they are cointegrated} \end{cases}$$

## Example: cointegration

Theory of purchasing power parity (PPP): "Apart from transportation costs, goods should sell for the same effective price in two countries"

$$P_t = S_t P_t^*$$

- $P_t$  is the price in USA (US \$)
- $P_t^*$  is the price in Denmark (DKK)
- $S_t$  is the exchange rate

Taking the log

$$p_t = s_t + p_t^*$$

## Example: cointegration

Usually that equation does not hold: measurement errors, transportation costs and differences in quality ( $z_t$ )

$$p_t = s_t + p_t^* + z_t$$

A weaker theory says that  $z_t \sim I(0)$  even though  $p_t, s_t, p_t^* \sim I(1)$ .  
These means that they are cointegrated.



# VAR in cointegration

Example of cointegrated system:

$$y_{1t} = \gamma y_{2t} + u_{1t}$$

$$y_{2t} = y_{2,t-1} + u_{2t}$$

Can this be written as a VAR for the differenced data?

$$\begin{bmatrix} \Delta y_{1t} \\ \Delta y_{2t} \end{bmatrix} = \begin{bmatrix} 1 - L & \gamma L \\ 0 & 1 \end{bmatrix} \begin{bmatrix} u_{1t} \\ u_{2t} \end{bmatrix}$$

Find the mistake in the equation above.

# VAR in cointegration

However, we can rewrite the VAR in levels as:

$$\begin{bmatrix} \Delta y_{1t} \\ \Delta y_{2t} \end{bmatrix} = \begin{bmatrix} -1 & \gamma \\ 0 & 0 \end{bmatrix} \begin{bmatrix} y_{1,t-1} \\ y_{2,t-1} \end{bmatrix} + \begin{bmatrix} u_{1t} + \gamma u_{2t} \\ u_{2t} \end{bmatrix}$$

With cointegrated systems, we have to include lagged levels along with lagged differences to explain  $\Delta \mathbf{y}_t$

# Cointegrating vector

Let  $\mathbf{y}_t = \{y_{1t}, y_{2t}\}'$  be a cointegrated vector of order one which components are composed by a common  $I(1)$  part and different  $I(0)$  parts.

$$\mathbf{y}_t = \begin{pmatrix} y_{1t} \\ y_{2t} \end{pmatrix} = (a, b) \begin{pmatrix} w_t \\ w_t \end{pmatrix} + \begin{pmatrix} \tilde{y}_{1t} \\ \tilde{y}_{2t} \end{pmatrix}$$

$$y_{1t} = aw_t + \tilde{y}_{1t}$$

$$w_t \sim I(1), \tilde{y}_{1t} \sim I(0)$$

$$y_{2t} = bw_t + \tilde{y}_{2t}$$

$$\tilde{y}_{2t} \sim I(0)$$

Then, what is

- $\beta(a, b)\mathbf{y}_t$ ?

# Cointegrating vector

- So there are infinite number of cointegrating vectors.
- We find the normalised one, the one that has a 1 as first element
- In our example above  $(1, a/b)$
- If  $\overbrace{\mathbf{y}_t}^{n \times 1} = (\mathbf{y}_{1,t}, \dots, \mathbf{y}_{n,t})$  how many cointegration vectors can we have?
- There might be  $h < n$  cointegrating vectors  $(n \times 1)$   $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_h$  such that  $\mathbf{a}_i' \mathbf{y}_t \sim I(0)$
- This means that there are  $n - h$  *common  $I(1)$  stochastic trends*
- All  $\mathbf{a}_i$  linearly independent (there is no scalar  $b$  such that  $\mathbf{a}_i = b\mathbf{a}_j$  for  $i \neq j$ )

# Cointegrating vectors

- $\mathbf{A}'\mathbf{y}_t \sim I(0)$  vector

$$\underbrace{\mathbf{A}'}_{h \times n} = \begin{bmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \\ \vdots \\ \mathbf{a}'_h \end{bmatrix}$$

- These vectors  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_h$  are not unique,
- So if  $\mathbf{A}'\mathbf{y}_t \sim I(0)$ , then  $\mathbf{b}'\mathbf{A}'\mathbf{y}_t \sim I(0)$  for  $\underbrace{\mathbf{b}'}_{1 \times h}$  any nonzero vector.
- Therefore the vector  $\boldsymbol{\pi} = \mathbf{b}'\mathbf{A}'$  could be considered as a cointegrating vector

# Simulating Cointegrated Systems

These systems can be simulated using the Phillip's triangular representation (Phillips, 1991)

Bivariate: 1 cointegrated vector, 1 common trend

- Assume  $\mathbf{y}_t = (y_{1t}, y_{2t})'$  is cointegrated,
- and  $\mathbf{a} = (1, -a_2)'$  is the cointegrating vector

$$y_{1t} = a_2 y_{2t} + e_{1t}$$

$$e_{1t} \sim I(0)$$

$$y_{2t} = y_{2,t-1} + e_{2t}$$

$$e_{2t} \sim I(0)$$

- First eq. describes the long-run equilibrium relationship with  $I(0)$  disequilibrium error  $e_{1t}$
- Second, specifies  $y_{2t}$  as the common stochastic trend with innovation  $e_{2t}$

## Example: Phillip's triangular representation

$$T = 250$$

$$\mathbf{a} = (1, -1)$$


$$e_{1t} = 0.75e_{1,t-1} + \nu_t$$

$$\nu_t \sim N(0, 0.5^2)$$

$$e_{2t} \sim IIDN(0, 0.5^2)$$

```
> library(splus2R)
> T=250
> e= rmvnorm(T, mean=rep(0,2), sd=c(0.5,0.5))
> e1.ar1= arima.sim(n=T,model=list(ar=0.75), innov= e[,1])
> y2= cumsum(e[,2])
> y1= y2 + e1.ar1
```

## Example: Phillip's triangular representation



They follow each other closely because their stochastic trend is the same.



# Simulating Cointegrated Systems

Trivariate: 1 cointegrated vector, 2 common trends

- Assume  $\mathbf{y}_t = (y_{1t}, y_{2t}, y_{3t})'$  is cointegrated,
- and  $\mathbf{a} = (1, -a_2, -b_3)'$  is the cointegrating vector

$$y_{1t} = a_2 y_{2t} + a_3 y_{3t} + e_{1t}$$

$$e_{1t} \sim I(0)$$

$$y_{2t} = y_{2,t-1} + e_{2t}$$

$$e_{2t} \sim I(0)$$

$$y_{3t} = y_{3,t-1} + e_{3t}$$

$$e_{3t} \sim I(0)$$

- First eq describes the long-run equilibrium relationship
- Second and third describe the common stochastic trends

## Example: Trivariate

$$T = 250$$

$$e_{1t} = 0.75e_{1,t-1} + \nu_t$$


$$e_{2t} \sim IIDN(0, 0.5^2)$$

$$\mathbf{a} = (1, -0.5, -0.5)$$

$$\nu_t \sim N(0, 0.5^2)$$

$$e_{3t} \sim IIDN(0, 4)$$

## Example: Trivariate



$y_2, y_3$  are two independent common trends,  $y_1$  is the average of the two trends plus a AR(1) residuals.

# Simulating Cointegrated Systems

Trivariate: 2 cointegrated vectors, 1 common stochastic trend

- Assume  $\mathbf{y}_t = (y_{1t}, y_{2t}, y_{3t})'$  is cointegrated,
- and  $\mathbf{a}_1 = (1, 0, -a_{13})'$  and  $\mathbf{a}_2 = (0, 1, -a_{23})$  cointegrated vectors

$$y_{1t} = a_{13}y_{3t} + e_{1t} \qquad e_{1t} \sim I(0)$$

$$y_{2t} = a_{23}y_{3,t-1} + e_{2t} \qquad e_{2t} \sim I(0)$$

$$y_{3t} = y_{3,t-1} + e_{3t} \qquad e_{3t} \sim I(0)$$

- First two eqs describe the two long-run equilibrium relationships
- Third describes the common stochastic trend

## Example: Trivariate

$$T = 250$$

$$\mathbf{a1} = (1, 0, -1) \quad \mathbf{a2} = (0, 1, -1)$$

$$e_{1t} = 0.75e_{1,t-1} + \nu_t$$


$$\nu_t \sim N(0, 0.5^2)$$

$$e_{2t} = 0.75e_{2,t-1} + \xi_t$$

$$\xi_t \sim N(0, 0.5^2)$$

$$e_{3t} \sim IIDN(0, 0.5^2)$$

# Example: Trivariate



# Error-correction representation (ECM)

For a bivariate VAR(p),

- Assume  $\mathbf{y}_t = (y_{1t}, y_{2t})'$  is cointegrated,
- and  $\mathbf{a} = (1, -a_2)'$  is the cointegrating vector

$$\Rightarrow \quad \mathbf{a}'\mathbf{y}_t = y_{1t} - a_2 y_{2t} \quad \text{is } I(0)$$

- Engle and Granger (1987) show that cointegration implies the existence of *error correction model* (ECM) describing the dynamic behaviour of  $y_{1t}$  and  $y_{2t}$ .

# Error-correction representation (ECM)

$$\begin{aligned}\Delta y_{1t} = & \alpha_1 + \pi_1 \underbrace{(y_{1,t-1} - a_2 y_{2,t-1})}_{\mathbf{a}'\mathbf{y}_t} \\ & + \underbrace{\sum_j \xi_{11}^j \Delta y_{1,t-j} + \sum_j \xi_{12}^j \Delta y_{2,t-j}}_{\text{short run dynamic adjustment}} + \epsilon_{2t}\end{aligned}$$

$$\begin{aligned}\Delta y_{2t} = & \alpha_2 + \pi_2 (y_{1,t-1} - a_2 y_{2,t-1}) \\ & + \sum_j \xi_{21}^j \Delta y_{1,t-j} + \sum_j \xi_{22}^j \Delta y_{2,t-j} + \epsilon_{2t}\end{aligned}$$



# Error Correction Model (ECM)

Intuition:  $z_t = y_{t1} - a_2 y_{2t} \sim I(0)$ , then  $y_{t1} - a_2 y_{2t}$  is the disequilibrium error

- Where the system goes at time  $t + 1$  depends on the sign and magnitude of the disequilibrium error at time  $t$ .
- Short-run dynamics are movements in the short run, modeled in the ECM, that guide the economy towards the long-run equilibrium  $y_{1t} = a_2 y_{2t}$

## Example: bivariate ECM for stock prices and dividends

- Let  $S_t$  be the stock price and  $D_t$  the annual dividend
- The dividend yield  $D_t/S_t$ , after taking logarithms become  $d_t - s_t$
- $\mathbf{y}_t = (s_t, d_t)'$  is  $I(1)$ .
- However the dividend yield  $d_t - s_t \sim I(0)$
- $\mathbf{a}' = (1, -1)$
- The theory says that there is a long-run equilibrium in the dividend yield:

$$d_t = s_t + \mu + e_t \quad e_t \sim I(0)$$

where  $\mu$  is the mean of the log dividend-price ratio and  $e_t$  is an  $I(0)$  random variable.

# Example: bivariate ECM for stock prices and dividends

The ECM has the form:

$$\Delta s_t = \alpha_s + \pi_s (d_{t-1} - s_{t-1} - \mu) + \epsilon_{st}$$

$$\Delta d_t = \alpha_d + \pi_d (d_{t-1} - s_{t-1} - \mu) + \epsilon_{dt}$$

where  $\alpha_s, \alpha_d > 0$ .

- The first equation relates the growth rate of stock prices to the lagged disequilibrium error  $d_{t-1} - s_{t-1} - \mu$
- The second eq. relates the growth rate of dividends to the lagged of the disequilibrium error
- The reactions to the disequilibrium error are adjusted by  $\pi_s, \pi_d$ .

# Example: bivariate ECM for stock prices and dividends

Let us assume that  $\pi_d = 0$  and  $\pi_s = 0.5$  (only  $s_t$  reacts to the disequilibrium)

- Case 1**  $d_{t-1} - s_{t-1} - \mu = 0$ , then  $E(\log s_t | \mathbf{y}_{t-1}) = \alpha_s$  and  $E(\log d_t) = \alpha_d$ . These quantities represent the growth rates of stock prices and dividends in long-run equilibrium.
- Case 2**  $d_{t-1} - s_{t-1} - \mu > 0$ , then  $E(\log s_t | \mathbf{y}_{t-1}) > \alpha_s$ . The dividend yield has increased above the long-run mean. The ECM predicts that  $s_t$  will grow faster than the equilibrium rate to restore the dividend yield to its long-run mean.
- Case 3**  $d_{t-1} - s_{t-1} - \mu < 0$ . The dividend yield has decreased below its long-run mean. ????

# Residual-based tests for cointegration

Let  $\mathbf{y}_t$  be a  $I(1)$  cointegrated vector:  $\mathbf{A}'\mathbf{y}_t \sim I(0)$

Testing for cointegration can be seen as testing for the existence of a long-run equilibria in the elements of  $\mathbf{y}_t$ . Two scenarios:

- ① There is at most one cointegrating vector (Engle and Granger, 1986)
  - Two-step residual-based test
- ② There are possibly  $0 \leq h < n$  cointegrating vectors (Johansen, 1988)
  - A more complicated procedure to determine the number of cointegrating relationships

# Engle and Granger's cointegration test

Once the cointegrating vector is pre-defined, we can test whether the residuals  $z_t = \mathbf{a}'\mathbf{y}_t$  are stationary.

- 1 Test whether each individual element of  $\mathbf{y}_t$  is  $I(1)$  – DF test or tests of stationarity
- 2 We construct a scalar  $z_t = \mathbf{a}'\mathbf{y}_t$  and we test  $H_0 : z_t \sim I(1)$  vs  $H_1 : z_t \sim I(0)$ .

## Conclusion:

If  $z_t$  is stationary then  $\mathbf{y}_t$  is cointegrated with cointegrating vector  $\mathbf{a}'$

## Example: PPP

Data from 1973:1 – 1989:10.


- $p_t = 100(\log(P_t) - \log(P_1))$  US consumer price
- $p_t^*$  Italy consumer price
- $s_t = -100(\log(S_t) - \log(S_1))$  dollar–lira exchange rate
- $z_t = p_t - s_t - p_t^*$

# Example: PPP

```
> load("../data/ppp.rda")
> selection <- window( ppp, start=c(1973,1), end=c(1989,10) )
> ppp.data <- cbind(
+   pstar=100*log(selection[, "PC6IT"]/selection[[1, "PC6IT"]]),
+   p=100*log(selection[, "PZUNEW"]/selection[[1, "PZUNEW"]]),
+   s=-100*log(selection[, "EXRITL"]/selection[[1, "EXRITL"]])
+ )
> ppp.data <- cbind( ppp.data,
+                   z = ppp.data[, "p"] - ppp.data[, "s"] - ppp.data[, "pstar"] )
```



# Example: PPP



## Example: PPP

Perform DF-test for the three series:

- $E(\Delta p_t) > 0$  so  $H_0 : p_t$  unit root process with positive drift vs  $H_1 : p_t$  stationary around a deterministic trend

```
> library(urca)
> s.df<-ur.df(as.vector(ppp.data[, "s"]), type="trend", lags=12)
> p.df<-ur.df(as.vector(ppp.data[, "p"]), type="trend", lags=12)
> z.df<-ur.df(as.vector(ppp.data[, "z"]), type="drift", lags=12)
```

## Example: PPP

```
> pstar.df<-ur.df(as.vector(ppp.data[, "pstar"]),  
+                 type="trend", lags=12)  
> attr(pstar.df, "teststat")
```

	tau3	phi2	phi3
statistic	-0.1319633	3.59166	4.249956

```
> attr(p.df, "teststat")
```

	tau3	phi2	phi3
statistic	-1.954675	2.066691	2.412933

```
> attr(s.df, "teststat")
```

	tau3	phi2	phi3
statistic	-1.584433	1.219635	1.489674

## Example: PPP

They are all individually  $I(1)$ . Theory says that  $\mathbf{a}' = (1, -1, -1)$ , then

$$z_t = p_t - s_t - p_t^*$$

The trends should be eliminated with this transformation, so we perform the ADF test on  $z_t$  with drift and get a statistics  $-2.04 > -2.88$  and so the null hypothesis of unit root is accepted.

The series are not cointegrated. At least not with this cointegrating vector

# Estimating the cointegrating vector

- If there is no theoretical candidate for  $\mathbf{a}$ , then it has to be estimated
- We find first a candidate with OLS
- If  $z_t = \mathbf{a}'\mathbf{y}_t$  is second moment stationary and ergodic:

$$\frac{1}{T} \sum_t z_t^2 = \frac{1}{T} \sum_t (\mathbf{a}'\mathbf{y}_t)^2 \rightarrow^p E(z_t^2)$$

- If instead,  $\mathbf{a}$  is not the cointegrating vector  $\Rightarrow z_t \sim I(1)$  and  $\frac{1}{T} \sum_t z_t^2 \rightarrow \infty$  as  $T \rightarrow \infty$
- So we find and estimate of the cointegrating vector by minimising  $\frac{1}{T} \sum_t (\mathbf{a}'\mathbf{y}_t)^2$  with respect to  $\mathbf{a}$

# Estimating the cointegrating vector

If we know for sure that  $a_1 \neq 0$ , then we set it to  $\hat{\mathbf{a}} = (1, -\hat{a}_2, -\hat{a}_3, \dots, -\hat{a}_n)$  result of the OLS estimation of:

$$y_{1t} = \alpha + a_2 y_{2t} + a_3 y_{3t} + \dots + a_n y_{nt} + \epsilon_t$$

- These estimates are super consistent (converges at rate  $T$ ), even if there is endogeneity.
- Their asymptotic distribution is non-normal.
- It might be substantially biased in small samples.
- A better estimator can be found.

# Estimating the cointegrating vector

$$y_{1t} = \alpha + a_2 y_{2t} + a_3 y_{3t} + \dots + a_n y_{nt} + \epsilon_t$$

- Let us estimate the cointegrating vector with OLS, what occurs if there is no cointegration relation? What does this estimate mean?
- $y_{1t} = \alpha + \mathbf{a}'_2 \mathbf{y}_{2t} + \epsilon_t$
- $\hat{\epsilon}_t = y_{1t} - \hat{\alpha} - \hat{\mathbf{a}}'_2 \mathbf{y}_{2t}$
- The unit root test on  $\hat{\epsilon}_t$  is without drift or trend
- All the elements in the regression are  $I(1)$ . So the OLS estimates will probably be spurious and  $\hat{\epsilon}_t \sim I(1)$
- So the distribution of the test have other type of asymptotic distributions known as the Phillips-Ouliaris (PO)
- These distributions also depend on whether  $y_{1t}, \mathbf{y}_{2t}$  have drift or not.

## Example: PPP

We say that  $p_t, p_t^*, s_t$  for the previous example were not cointegrated with cointegrating vector  $(1, -1, -1)$ . What about for another linear relationship?

```
> ppp.lm<-lm( p ~ 1 + s + pstar, ppp.data )
> epsilon.hat<-resid(ppp.lm)
> epsilon.hat.df<-ur.df(epsilon.hat, type="none", lags=12)
> attr(epsilon.hat.df, "teststat")
```

```
tau1
statistic -2.73094
```

$$p_t = 2.71 + 0.05s_t + 0.53p_t^* + \hat{\epsilon}_t$$

We are in a case 3 of test for cointegration (Table 19.1). The critical values for the DF test are in Table B.9. There is little evidence of cointegration



## Efficient lead/lag estimator

- Stock and Watson (1993) amongst others suggest an asymptotically efficient estimator (equivalent to MLE) for normalised cointegrating vector  $\mathbf{a} = (1 - \mathbf{a}_2)$
- Estimate the following augmented the cointegrating regression by OLS

$$y_{1t} = (\alpha + \delta t) + \mathbf{a}_2 \mathbf{y}_{2t} + \sum_{j=-p}^p \rho_j \Delta \mathbf{y}_{2,t-j} + u_t$$

- The resulted  $\hat{\mathbf{a}}_2$  is called the *dynamic OLS* (DOLS) estimator which is consistent, asymptotically normal and efficient under certain conditions
- Asymptotically valid standard errors are given by the OLS standard errors multiplied by a ratio

# Example: Income/consumption